

HYDROBROMOCARBON BLENDS TO PROTECT AGAINST FIRES AND EXPLOSIONS

Publication number: WO9809686

Publication date: 1998-03-12

Inventor: TAPSCOTT ROBERT E (US); MOORE TED A (US); LIFKE JOSEPH L (US)

Applicant: UNIV NEW MEXICO (US); TAPSCOTT ROBERT E (US); MOORE TED A (US); LIFKE JOSEPH L (US)

Classification:






- international: **A62C35/02; A62D1/00; A62D1/08; A62C35/00; A62D1/00; (IPC1-7): A62D**

- european: A62D1/00C6

Application number: WO1997US16660 19970909






Priority number(s): US19960025969P 19960909

Also published as:

 WO9809686 (A3)
 WO9809686 (A3)
 EP0930918 (A3)
 EP0930918 (A3)
 EP0930918 (A2)

more >>

Cited documents:

 US5219474
 US2569979
 US2818382
 US3657120
 US5124053

more >>

Report a data error here

Abstract of **WO9809686**

A set of blends for extinguishment of fires, suppression of explosions, and inertion against fires and explosions is disclosed. The blends are comprised of a bromine-containing component and a fluorine-containing component. The bromine-containing component is comprised of one or more hydrobromoalkanes, hydrobromoalkenes, and/or hydrobromoarenes. The fluorine-containing component is comprised of one or more fluorine-containing halocarbons, which may also contain chlorine. Specifically provided as fluorine-containing halocarbons are saturated and unsaturated hydrochlorofluorocarbons, hydrofluorocarbons, perfluorocarbons, perfluoroethers, hydrofluoroethers, and/or hydrofluoropolyethers. This provides a blend that mimics the fire and explosion protection action of halons and hydrobromofluorocarbons without the environmental impact associated with these compounds, which contain bromine and fluorine in the same molecule.

Data supplied from the **esp@cenet** database - Worldwide



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A62D	A2	(11) International Publication Number: WO 98/09686 (43) International Publication Date: 12 March 1998 (12.03.98)
(21) International Application Number: PCT/US97/16660 (22) International Filing Date: 9 September 1997 (09.09.97) (30) Priority Data: 60/025,969 9 September 1996 (09.09.96) US (71) Applicant (for all designated States except US): THE UNIVERSITY OF NEW MEXICO [US/US]; Albuquerque, NM 87131 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): TAPSCOTT, Robert, E. [-/US]; 3812 Palomas Drive, N.E., Albuquerque, NM 87110 (US). MOORE, Ted, A. [-/US]; 6300 Acacia Street, N.M., Albuquerque, NM 87120 (US). LIFKE, Joseph, L. [-/US]; 2900 Vista Del Rey, N.E., #15B, Albuquerque, NM 87112 (US). (74) Agent: ROBERT W. BECKER & ASSOCIATES; Suite B, 11896 N. Highway 14, Tijeras, NM 87059 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: HYDROBROMOCARBON BLENDS TO PROTECT AGAINST FIRES AND EXPLOSIONS (57) Abstract <p>A set of blends for extinguishment of fires, suppression of explosions, and inertion against fires and explosions is disclosed. The blends are comprised of a bromine-containing component and a fluorine-containing component. The bromine-containing component is comprised of one or more hydrobromoalkanes, hydrobromoalkenes, and/or hydrobromoarenes. The fluorine-containing component is comprised of one or more fluorine-containing halocarbons, which may also contain chlorine. Specifically provided as fluorine-containing halocarbons are saturated and unsaturated hydrochlorofluorocarbons, hydrofluorocarbons, perfluorocarbons, perfluoroethers, hydrofluoroethers, and/or hydrofluoropolyethers. This provides a blend that mimics the fire and explosion protection action of halons and hydrobromofluorocarbons without the environmental impact associated with these compounds, which contain bromine and fluorine in the same molecule.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

HYDROBROMOCARBON BLENDS TO PROTECT AGAINST FIRES AND EXPLOSIONS

Government Rights

This invention was made under contract with the U.S.
Government, which has certain rights therein.

Field of the Invention

The invention described and claimed herein is generally related to chemical agents used for fire extinguishment, explosion suppression, explosion inertion, and fire inertion, and more particularly, to extinguishing, suppressing, and inerting blends of hydrobromoalkanes, hydrobromoalkenes, and hydrobromoarenes with fluorine-containing halocarbons to provide replacements for halon fire and explosion suppressants and extinguishants. The production of halons has been eliminated or curtailed due to their impact on stratospheric ozone.

Background of the Invention and Prior Art

The broad class of compounds termed "halocarbons" consists of all molecules containing carbon (C) and one or more of the atoms fluorine (F), chlorine (Cl), bromine (Br), and/or iodine (I). These four elements—fluorine, chlorine, bromine, and iodine—are members of the halogen family of elements. When defined in the broadest sense, as we do here, halocarbons may also contain other chemical features such as hydrogen atoms, carbon-to-carbon multiple bonds, aromatic rings, and ether linkages. Haloalkanes, a subset of halocarbons, contain only single bonds between the carbon atoms. Haloalkenes contain one or more double bonds connecting carbon atoms. Haloarenes contain aromatic groups based on the six-carbon benzene ring. Aromatic groups formally contain alternating single and double carbon to carbon bonds, but in actuality, the bonds are "delocalized" such that the carbon to carbon bonding is greater than single bonding but less than double bonding. Compounds that contain no multiple bonding, such as the alkanes, are said to be "saturated." Alkenes and arenes are said to be "unsaturated."

The use of certain haloalkanes as fire extinguishing agents has been known for many years. Fire extinguishers containing carbon tetrachloride (CCl_4 , also known as tetrachloromethane) or methyl bromide (CH_3Br , also known as bromomethane) were used in aircraft applications as early as the 1920s (Charles L. Ford, "An Overview of Halon 1301 Systems," in Halogenated Fire Suppressants, Richard G. Gann, editor, ACS Symposium Series 16, American Chemical Society, Washington, DC, 1975). Over a period of years, the high toxicity of these compounds was recognized and they were replaced with less toxic materials, in particular bromofluoroalkanes and closely related compounds. A major study of haloalkanes as fire extinguishing agents was conducted by the Purdue Research Foundation for the U.S. Army from 1947 to 1950 (Fire Extinguishing Agents, Final Report, Purdue University, 1950. A synopsis of the results from this study is available in Ford, op. cit.).

The term "extinguishment" is usually used to denote complete elimination of a fire; whereas, "suppression" is often used to denote reduction, but not necessarily total elimination, of a fire or explosion. These two terms are sometimes used interchangeably. There are four general types of halocarbon fire and explosion protection applications. (1) In total-flood fire extinguishment and/or suppression applications, the agent is discharged into a space to achieve a concentration sufficient to extinguish or suppress an existing fire. This is often, though not always, done by an automatic system, which detects the fire and then automatically discharges the extinguishing agent to fill the space with the concentration of a gaseous or an evaporated volatile liquid agent to the concentration needed to suppress or extinguish the contained fire. Total flooding use includes protection of enclosed, potentially occupied spaces such as computer rooms as well as specialized, often unoccupied spaces such as aircraft engine nacelles and engine compartments in vehicles. Note that the term "total flood" does not necessarily mean that the extinguishing or suppressing agent is uniformly dispersed

throughout the space protected.³ (2) In streaming applications, the agent is applied directly onto a fire or into the region of a fire. This is usually accomplished using manually operated wheeled or portable fire extinguishers. A second method, which we have chosen to include as a streaming application, uses a "localized" system, which discharges agent toward a fire from one or more fixed nozzles. Localized systems may be activated either manually or automatically. (3) In explosion suppression, an agent is discharged to suppress an explosion that has already been initiated. The term "suppression" is normally used in this application since an explosion is usually self-limiting. However, the use of this term does not necessarily imply that the explosion is not extinguished by the agent. In this application, a detector is usually used to detect an expanding fireball from an explosion, and the agent is discharged rapidly to suppress the explosion. Explosion suppression is used primarily, but not solely, in military applications. (4) In inertion, an agent is discharged into a space to prevent an explosion or a fire from being initiated. Often, a system similar or identical to that used for total-flood fire extinguishment or suppression is used. Inertion is widely used for protection of oil production facilities at the North Slope of Alaska and in other areas where flammable gases or explosive dusts may build up. Usually, the presence of a dangerous condition (for example, dangerous concentrations of flammable or explosive gases) is detected, and the agent is then discharged to prevent the explosion or fire from occurring until the condition can be remedied.

Thus, there are four fire and explosion protection applications covered by this disclosure:

1. Total-Flood Fire Extinguishment and Suppression
2. Streaming Fire Extinguishment and Suppression
3. Explosion Suppression
4. Explosion and Fire Inertion

The cup burner is a widely accepted laboratory test apparatus for determining the fire extinguishing and suppressing effectiveness of

4

agents. In this method, an agent is introduced into a stream of air which passes around a cup of burning liquid fuel, and the concentration of gaseous agent needed to extinguish the flame is determined. During this operation, any agent that is normally a liquid is allowed to become a gas before being mixed into the stream of air and passed by the burning liquid fuel. The cup burner is so widely accepted that the National Fire Protection Association (NFPA) Standard 2001 on Clean Agent Fire Extinguishing Systems mandates this method as the primary procedure for determining the concentration needed to extinguish a fire of liquid hydrocarbon fuels (e.g., gasoline, hexane, etc.; such fires are termed "Class B fires"). That standard states that "The minimum design concentration for Class B flammable liquids shall be a demonstrated extinguishing concentration plus a 20 percent safety factor. Extinguishing concentration shall be demonstrated by the cup burner test." Concentrations are usually expressed as "percent by volume." This is the same as the "percent by gas volume," which is calculated assuming that all of the introduced agent has volatilized (i.e., vaporized to become a gas).

The halocarbons most widely used for fire extinguishment (by total flooding or streaming), explosion suppression, explosion inertion, and fire inertion have been the three compounds shown in Table I. These materials are all alkanes containing both bromine and fluorine. To avoid the use of complicated chemical names, these (and other halocarbons used for fire and explosion protection) are often designated by a "Halon Number." Usually the word "Halon" is capitalized when used as part of a halon number, but is not capitalized when used generically for haloalkanes employed in fire and explosion protection. In recent years and, in particular, in regulatory documents, the term "halon" has been increasingly applied to denote the specific, widely used halocarbon agents shown in Table I and this is a practice that we use here. The "CAS No." is the number assigned by the Chemical Abstract Services of the American Chemical Society to aid in identifying chemical compounds. Halon 1301 has been widely used for

total-flood fire extinguishment, explosion suppression, and inertion. Due to its higher boiling point and higher toxicity, Halon 1211 is most often used in streaming. Halon 2402 has had significant use in Eastern Europe for both total-flood and streaming, but has had relatively little use in other parts of the world.

TABLE I. HALONS USED IN FIRE AND EXPLOSION PROTECTION

Name	Formula	Halon No.	CAS No.	ODP
bromotrifluoromethane	CBrF_3	1301	75-63-8	10
bromochlorodifluoromethane	CBrClF_2	1211	353-59-3	3
1,2-dibromotetrafluoroethane	$\text{CBrF}_2\text{CBrF}_2$	2402	124-73-2	6

Bromine-containing compounds such as the halons are believed to operate as fire extinguishing agents by a complex chemical reaction mechanism involving the disruption of free-radical chain reactions, which are essential for continued combustion. Bromine is much more effective than chlorine or fluorine in promoting this disruption. In fact, there is doubt that chlorine or, in particular, fluorine plays a significant role in free-radical reaction disruption. The fluorine-containing portion of the halon molecules may, however, provide significant cooling and may thereby enhance extinguishment by the bromine. The halons are desirable as fire extinguishing agents because they are effective, because they leave no residue (i.e., they are liquids that evaporate completely or they are gases), and because they do not damage equipment or facilities to which they are applied.

Recently however, the halons have come to be recognized as serious environmental threats due to their ability to cause stratospheric ozone depletion. The ability of a compound to deplete stratospheric ozone is termed the "Ozone Depletion Potential" (ODP). Larger ODPs indicate greater stratospheric ozone depletion. ODPs reported for the

halons are shown in Table I (Federal Register, Vol. 58, No. 250, 30 December 1991). Since ODPs are calculated and various models and input data are used, other values have been reported. The relatively high ODPs of the halons is due to two factors. (1) Bromine is a potent depleter of stratospheric ozone and is much more damaging to ozone than is chlorine. (2) Fluorine increases compound stability, greatly decreasing the amount of compound removal and breakdown in the troposphere and thus allowing most of any discharged halon to reach the stratosphere, where ozone destruction occurs. Due to stratospheric ozone depletion concerns, the Montreal Protocol, an international treaty prepared in 1987 and amended several times since, has required a halt in the production of Halon 1301, Halon 1211, and Halon 2402 at the end of 1993 in the United States and in other industrialized nations.

Much research has gone on to find replacements for the halons for protection against fires and explosions; however, the search for halon replacements has been less than totally successful ("Pressure Mounts As Search for Halon Replacements Reaches Critical Phase," Chemical and Engineering News, September 19, 1994, pp. 29-32). One class of candidates proposed at one time as replacements of the halons were the hydrobromofluorocarbons (HBFCs), which contain only hydrogen, bromine, fluorine, and carbon. For example, bromodifluoromethane, CHBrF_2 , is a highly effective HBFC fire suppressant and was commercialized for a short time as a halon replacement. Like the halons, however, the production of most HBFCs has now been banned in industrialized nations due to their significant ODP. None of the agents now being actively promoted as replacements for the halons contain bromine. The primary replacements are now hydrochlorofluorocarbons (HCFCs), hydrofluorocarbons (HFCs), and perfluorocarbons (PFCs or FCs). HCFCs, HFCs, and PFCs (FCs) appear to operate primarily by heat absorption, which is a less effective mechanism for most fire and explosion protection applications than the free-radical chain disruption mechanism believed to be effected by

7

bromine and believed to be the primary mechanism for fire extinguishment by the halons. Thus, HCFCs, HFCs, and PFCs (a group of materials that we refer to as "first-generation" halon replacements) have a significantly decreased effectiveness in most fire and explosion protection applications compared to the halons that they are replacing.

Although bromine is believed to be the primary feature providing the outstanding fire extinguishment capability of the halons, it is precisely this feature that causes most (for Halon 1301 and Halon 2402, essentially all) of the stratospheric ozone depletion exhibited by these agents. Similarly, although fluorine-containing portions of the halon molecules may provide significant cooling and thereby enhance fire suppression by the bromine, it is precisely this portion of the chemicals that stabilizes the molecule and allows the halons to reach the stratosphere, where ozone depletion occurs. Thus, it is the combination of bromine and fluorine that both provides the outstanding fire extinguishment and produces the large ozone depletion exhibited by halons. Note, however, that it is the combination of bromine and fluorine in the same molecule that leads to large ozone depletion impacts. We have conceived, therefore, that one should be able to use a blend of two or more materials, none of which contain bromine and fluorine in the same molecule, to produce at a fire a mixture of bromine- and fluorine-containing fragments and bromine and fluorine atoms such as would be produced by the halons and by the HBFCs without the associated environmental impacts and regulatory restrictions. We have thus conceived blends of a bromine-containing component comprised of one or more halocarbons containing only bromine as the halogen and a fluorine-containing component comprised of one or more halocarbons containing only fluorine or only fluorine and chlorine as the halogens.

We first conceptualized the hydrobromoalkanes (compounds containing only hydrogen, bromine, and carbon) as constituents of the bromine-containing component. Of particular importance is whether such compounds would have sufficiently low ODPs to make them

environmentally acceptable. The lightest member of the family of hydrobromoalkane chemicals, methyl bromide, has an unacceptably high ODP of 0.64 (Scientific Assessment of Ozone Depletion: 1994, Report No. 37, National Oceanic and Atmospheric Administration, National Aeronautics and Space Administration, United Nations Environment Programme, and World Meteorological Organization, February 1995) and is undergoing increasing restrictions due to ozone depletion concerns. We conceived, however, that hydrobromoalkanes with more hydrogen atoms and with more carbon-carbon bonds than methyl bromide should have lower ODPs and might not be environmentally harmful. For example, the atmospheric lifetimes of perfluorocarbons, which contain no hydrogen atoms, decreases from 50,000 years for carbon tetrafluoride (CF_4), which has no carbon-carbon bonds, to 10,000 years for hexafluoroethane (CF_3CF_3), with one carbon-carbon bond, to 2600 years for octafluoropropane ($\text{CF}_3\text{CF}_2\text{CF}_3$), with two carbon-carbon bonds (Climate Change 1995, The Science of Climate Change, J. T. Houghton, L. G. Meira Filho, B. A. Callander, N. Harris, A. Kattenberg, and K. Maskell, editors, Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, UK, 1996.). Compounds that are rapidly removed from the atmosphere and, therefore, have a decreased atmospheric lifetime, will have a decreased impact on stratospheric ozone. Methyl bromide is the only hydrobromoalkane for which an ODP (or atmospheric lifetime) has been reported. To determine whether our concept was correct, we first used reported hydroxyl reaction rate constants to calculate atmospheric lifetimes for several hydrocarbons with increasing numbers of hydrogen atoms and carbon-carbon bonds. The results are given in Table II. As we expected, the atmospheric lifetime decreases with increasing hydrogen content and carbon bonding. We then plotted the values in Table II and obtained a rather smooth curve (Figure 1). What was surprising and was unexpected was that the atmospheric lifetime value for bromomethane, the only hydrobromoalkane for which an

atmospheric lifetime has been reported, appears to lie on the extrapolated curve. We therefore postulated that hydrobromoalkanes having a single bromine atom, would all lie on or near the curve obtained for the simple hydrocarbons. For example, the 1-bromopropane, a hydrobromoalkane with seven hydrogen atoms and two carbon-carbon bonds, would be predicted to have an atmospheric lifetime of around 30 days. We have concluded from reported values for ODPs and atmospheric lifetimes that for bromine-containing compounds, the ODP increases by approximately 2 for each 10-year increase in atmospheric lifetime. Thus, with a predicted lifetime of about 1 month for bromopropane, one would predict an ODP of approximately 0.017. Bromobutanes containing a single bromine atom (e.g., 1-bromobutane, $\text{CH}_3\text{CHBrCH}_2\text{CH}_3$, and 2-bromobutane, $\text{CH}_3\text{CHBrCH}_2\text{CH}_3$), hydrobromoalkanes with nine hydrogen atoms and three carbon-carbon bonds, have predicted atmospheric lifetimes of approximately 7 days and ODP values of approximately 0.004. It should be noted that the atmospheric lifetimes may also decrease as the number of bromine atoms increases due to photolysis—decomposition by solar radiation, another, but often less efficient, atmospheric removal process. Thus, our investigation indicated that methyl bromide has an “anomalously” high ODP and that hydrobromoalkanes containing more hydrogen atoms and/or carbon-carbon bonds should be environmentally acceptable, as might such molecules with multiple bromine atoms.

10
TABLE II. TROPOSPHERIC HYDROXYL RATE CONSTANTS
AND LIFETIMES CALCULATED FROM THOSE CONSTANTS
FOR ALKANES.

IUPAC name	Formula	Number of hydrogen atoms	Atmospheric lifetime, days
methane	CH ₄	4	154
ethane	C ₂ H ₆	6	46.5
propane	C ₃ H ₈	8	9.49
n-butane	C ₄ H ₁₀	10	4.67

5

10 To determine whether our concept for a fire extinguishing agent was valid, we tested two blends of 1-bromopropane (CH₂BrCH₂CH₃, 10 percent and 25 percent by weight) as the bromine-containing component with a hydrofluoropolyether as the fluorine-containing component. The hydrofluoropolyether was a mixture of different

15 molecular weight materials containing ether (C-O-C) linkages and hydrogen and fluorine substituents. The materials were tested using a cup burner apparatus and *n*-heptane fuel (C₇H₁₆). For five determinations each, the average cup-burner extinguishment concentrations (with average deviations) were 3.18 ± 0.05 volume

20 percent in air for the 25 percent 1-bromopropane blend, 3.11 ± 0.04 volume percent in air for the 10 percent blend, 5.23 ± 0.10 volume percent in air for the hydrofluoropolyether by itself, and 4.63 ± 0.23 volume percent in air for the 1-bromopropane by itself. A lower volume percent in air required for extinguishment indicates better

25 performance. The results were surprising for three reasons. First, the blends were better than either of the two components separately. While we had hoped that this would be true, the magnitude of the improvement was unexpected. The average extinguishment concentrations for the two blends were approximately 40 percent lower

30 than the extinguishment concentration for the hydrofluoropolyether by

itself and approximately 32 percent lower than the extinguishment concentration obtained with 1-bromopropane by itself. Second, the extinguishment concentrations exhibited by the blends were very close to those obtained in separate studies for Halon 1301 and Halon 1211 (approximately 2.9 and 3.2 percent, respectively). This was entirely unexpected since it has proven extremely difficult to find halon replacement candidates with extinguishment concentrations as low as the halons. In fact, these cup burner extinguishment concentrations are better than those reported for any agents now being commercialized (NFPA 2001 Standard on Clean Agent Fire Extinguishing Systems 1996 Edition, National Fire Protection Association, 1 Batterymarch Park, Quincy, Massachusetts, 1996). Third, although the difference was small and lies within the data scatter, the blend containing 10 percent 1-bromopropane appeared to be slightly more effective than the blend containing 25 percent 1-bromopropane. One would expect that the performance should improve as the bromine concentration increased.

Field testing also indicated the surprising result that a lower concentration of 1-bromopropane gave improved performance. Thus, in streaming tests, a blend of 25 percent 1-bromopropane with 75 percent hydrofluoropolyether required a flow rate of 0.29 pounds/second to extinguish a 2.25-square foot heptane fire, whereas a blend of 10 percent 1-bromopropane with 90 percent hydrofluoropolyether required a flow rate of only 0.17 pounds/second. In these tests, a lower flow rate for extinguishment implies an improved performance. A flow rate of 0.38 pounds/second was required for extinguishment with the hydrofluoropolyether by itself. Again the effectiveness of the blends and the difference between the 10 percent 1-bromopropane and 25 percent 1-bromopropane blends was surprising and entirely unexpected.

Because of these unexpectedly good results, we looked for other families of compounds resembling the bromoalkanes in that they

12

were composed only of bromine, hydrogen, and carbon. Hydrobromoalkenes and hydrobromoarenes meet the requirements and, moreover, have atmospheric lifetimes and ODPs even lower than those of the hydrobromoalkanes. The presence of multiple bonding in these unsaturated compounds provides additional reaction pathways for removal from the atmosphere. Thus, hydrobromoalkenes and hydrobromoarenes are unlikely to be regulated. We, therefore, conceived that hydrobromoalkenes and hydrobromoarenes could also be used in the bromine-containing components of the conceptualized blends. Some fluorine-containing bromoalkenes have been suggested for investigation as fire suppressants (W. M. Pitts, M. R. Nyden, R. G. Gann, W. G. Mallard, and W. Tsang, Construction of an Exploratory List of Chemicals to Initiate The Search for Halon Alternatives, NIST Technical Note 1279, Air Force Engineering and Services Laboratory, Tyndall Air Force Base, Florida, National Institute of Standards and Technology, Gaithersburg, Maryland, August 1990). Nonblended fluorine-containing bromoalkenes and nonblended fluorine-containing bromoarenes have been proposed as fire extinguishants (R. E. Tapscott, G. D. Brabson, G. W., Gobeli, E. W., Heinonen, J. A., Kaizerman, J. L., Lifke, and R. A. Patterson, "Research on Advanced Agents as Halon Replacements," Proceedings, International Conference on Ozone Protection Technologies, Washington, D.C., pp. 651-658, 21-23 October 1996; M. L. Robin, "Halogenated Fire Suppression Agents," in Halon Replacements, Technology and Science, ACS Symposium Series 611, Miziolek, A. W. and Tsang, W., editors, American Chemical Society, Washington, DC, 9, pp. 85-98, 1995.) and cup burner flame extinguishment concentrations for some fluorinated bromoalkenes have been reported (Tapscott, R. E., and Mather, J. D., Development of a Tropodegradable Total-Flooding Agent, Phase II: Initial Screening, NMERI 96/22/30930, Advanced Agent Working Group, July 1997). In none of this art, however, were any alkenes or arenes containing no fluorine proposed as fire suppressants and no blends of these

13

compounds were proposed, probably because the thinking up to now has been that fluorine is needed in the same compound as bromine in order to have a workable fire suppressant

Although the purpose of the fluorine-containing component is to add fluorine to the blend reaching the fire to mimic the action of halons and HBFCs, there are side benefits. For example, the use of a nonflammable or a low-flammability fluorine-containing component may allow the use of normally flammable constituents in the bromine-containing component. Moreover, fluorine-containing components with appropriate physical properties may provide improvements in discharge and dispersion of bromine-containing materials having very high or very low boiling points. Compounds with very high boiling points may not disperse effectively to fill a space and compounds with very low boiling points may not discharge well in streaming applications. In addition, suitable fluorine-containing components can decrease toxicological concerns that may be associated with certain hydrobromocarbons by diluting the bromine-containing material. Our work also indicates that some blends possess flame extinguishment and suppression ability greater than would be predicted from the intrinsic fire suppression ability of the separate components, a phenomenon that we term "synergism."

Accordingly, it is an object of the present invention to provide effective fire extinguishing, fire suppression, explosion suppression, and explosion and fire inertion blends that contain two components.


Summary of the Invention

This object is realized by providing a bromine-containing component comprised of one or more hydrobromocarbons, specifically the hydrobromoalkanes, hydrobromoalkenes, and hydrobromoarenes and a fluorine-containing component is comprised of one or more fluorine-containing halocarbons that contain no bromine and also no iodine.


The present invention therefore provides blends of hydrobromocarbons (specifically, hydrobromoalkanes, hydrobromoalkenes, and hydrobromoarenes) with halocarbons that always contain fluorine and, in some cases, also chlorine (but no bromine or iodine) for use as agents for fire extinguishing and suppression (in either total-flooding or streaming application), explosion suppression, and explosion and fire inertion. Note that in this application, "Blend" and "mixture" are used interchangeably.

In particular, the blend can be disposed, for example, in a pressurized discharge system and is adapted to be discharged into an area, for example to provide an average resulting concentration in such area of between 1-15%, and preferable 3-10% by gas volume, to extinguish or suppress a fire in that area. To suppress an explosion, a gas volume of 1-40% and preferably 5-20% is desired, while to prevent a fire or explosion from occurring, 1-30% and preferably 3-12% by gas volume is desired.

As the term is used in this application, hydrobromoalkanes are any compounds containing one or more bromine atoms and one or more hydrogen atoms attached to a linear, branched, or cyclic carbon chain or a combination of such chains and having no double bonds. Examples of such chains are shown below. Specifically excluded are the single-carbon compounds bromomethane, CH_3Br , which is known to cause environmental concerns, and dibromomethane (CH_2Br_2) and tribromomethane (CHBr_3), which have no carbon-carbon bonds and are predicted to have sufficiently long atmospheric lifetimes that they will have unacceptable ODPs.




Linear




Branched

15




Cyclic




Combination

5


As the term is used in this application, hydrobromoalkenes are any compounds containing one or more bromine atoms and one or more hydrogen atoms attached to a linear, branched, or cyclic carbon chain or a combination of such chains having one or more double bonds. Examples of such chains are shown below.




Linear



Branched




Cyclic




Combination

10



Linear With Two Double Bonds




Cyclic With Two Double Bonds




15

As the term is used here, hydrobromoarenes contain bromine and hydrogen in a molecule that contains one or more "aromatic" rings of carbon atoms. The most common of these is the six-carbon benzene ring, which, formally, contains alternating single and double bonds. Actually, the double bonding is "delocalized" such that each bond is equivalent to 1-1/2 bonds. Rings can also be joined to form additional

aromatic compounds, and may contain alkyl groups. Alkyl groups are groups containing only carbon and hydrogen atoms such as methyl (-CH₃), ethyl (-CH₂CH₃), *n*-propyl (-CH₂CH₂CH₃), *iso*-propyl (-CH(CH₃)₂), and *cyclo*-butyl (-C₄H₇). The bromine atoms may be attached directly to the aromatic ring, to alkyl substituents, or to a combination of these. Examples of carbon chains in arenes, without the bromine or hydrogen substituents, are shown below.



Benzene Ring

Two Fused Benzene
Rings (Naphthalene Ring System)Two Attached Benzene Rings
(Biphenyl Ring System)

Alkyl-Substituted Benzene Ring

Hydrobromoalkanes

Hydrobromoalkanes include, by way of example only, the linear and branched monobromo compounds such as CH₃CH₂Br, CH₃CH₂CH₂Br, CH₃CH₂CH₂CH₂Br, CH₃CHBrCH₃, CH₃CH(CH₃)CH₂Br and, in general, compounds having a formula C_nH_{2n+1}Br, where "n" is 2 or greater. Also disclosed here are the linear and branched dibromo compounds such as CH₃CHBr₂, CH₂BrCH₂Br, CH₃CH₂CHBr₂, CH₃CHBrCH₂Br, CH₃CHBr(CH₃)CH₂Br and, in general, compounds having a formula C_nH_{2n}Br₂, where "n" is 2 or greater. In general, the formulas of all of the linear and branched hydrobromoalkanes disclosed here have the formula C_nH_{2n+2-x}Br_x, where

17


"n" is 2 or greater and "x" is at least 1, but not larger than 2n+1. A list of some linear and branched hydrobromoalkanes is shown in Table III.

TABLE III. SELECTED LINEAR AND BRANCHED
HYDROBROMOALKANES.

	Formula	Name
5	$\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$	1-bromopropane
	$\text{CH}_3\text{CHBrCH}_3$	2-bromopropane
	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Br}$	1-bromobutane
	$\text{CH}_3\text{CH}_2\text{CHBrCH}_3$	2-bromobutane
10	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{Br}$	1-bromopentane
	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CHBrCH}_3$	2-bromopentane
	$\text{CH}_3\text{CH}_2\text{CHBrCH}_2\text{CH}_3$	3-bromopentane
	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{Br}$	1-bromohexane
	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CHBrCH}_3$	2-bromohexane
15	$\text{CH}_3\text{CH}_2\text{CHBrCH}_2\text{CH}_2\text{CH}_3$	3-bromohexane
	$\text{CH}_3\text{CH}(\text{CH}_3)\text{CH}_2\text{Br}$	1-bromo-2-methylpropane
	$\text{CH}_3\text{C}(\text{CH}_3)\text{BrCH}_3$	2-bromo-2-methylpropane
	$\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{Br}$	1-bromo-2-methylbutane
	$\text{CH}_3\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{Br}$	1-bromo-3-methylbutane
20	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{Br}$	1-bromo-2-methylpentane
	$\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{Br}$	1-bromo-3-methylpentane
	$\text{CH}_3\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_2\text{Br}$	1-bromo-4-methylpentane
	CH_3CHBr_2	1,1-dibromoethane
	$\text{CH}_2\text{BrCH}_2\text{Br}$	1,2-dibromoethane
25	$\text{CH}_3\text{CH}_2\text{CHBr}_2$	1,1-dibromopropane
	$\text{CH}_3\text{CHBrCH}_2\text{Br}$	1,2-dibromopropane
	$\text{CH}_2\text{BrCH}_2\text{CH}_2\text{Br}$	1,3-dibromopropane


18


Hydrobromoalkanes also include cyclic compounds, which contain rings of carbon atoms. These include the cyclic monobromo compounds such as C_3H_5Br , C_4H_7Br , C_5H_9Br , $C_6H_{11}Br$, and, in general, cyclic compounds having a formula $C_nH_{2n-1}Br$; the cyclic dibromo compounds such as $C_3H_4Br_2$, $C_4H_6Br_2$, $C_5H_8Br_2$, $C_6H_{10}Br_2$, and, in general, cyclic compounds having a formula $C_nH_{2n-2}Br_2$; and more highly bromine-substituted cyclic bromocarbons. Cyclic hydrobromocarbons may also contain multiple rings. Thus, for example, a dibromo cyclic hydrobromoalkane containing two joined four-membered rings would have the formula $C_8H_{12}Br_2$. All of the cyclic hydrobromoalkanes disclosed here have the general formula $C_nH_{2n+2-2y-x}Br_x$, where "n" is 3 or greater, "x", is at least 1, but not larger than $2n+1-2y$, and y is the number of rings. In general, hydrobromoalkanes containing more than one bromine atom can exist in more than one isomeric form. Example structures are shown below for cyclic hydrobromoalkanes.



$C_4H_6Br_2$ (3 Isomers Possible)

Hydrobromoalkanes also include cyclic compounds with alkyl substituents such as those shown below.



C₇H₁₃Br₂ (Many Isomers Possible)

5 Hydrobromoalkenes

Hydrobromoalkenes include the linear and branched compounds containing one carbon-carbon double bond, one or more hydrogen atoms, and one or more bromine atoms. Examples are CH₂=CHBr, CH₂=CHCH₂Br, CH₂=CBrCH₃, CHBr=CHCH₃, CH₂=CBrCH₂Br, CHBr=C(CH₃)CH₃, CH₂=CHCHBr₂, CBr₂=C(CH₃)CH₂Br, and, in general, hydrobromoalkenes having a formula C_nH_{2n-x}Br_x, where "n" is 2 or greater and "x" is 2n-1 or less but not less than one. They also include the linear and branched compounds containing two carbon-carbon double bonds and one or more bromine atoms CH₂=CHCH=CHBr, CH₂=CHCBr=CH₂, CH₂=C(CH₃)CBr=CH₂, CH₂=C(CH₃)CBr=CHBr, and, in general, hydrobromoalkenes having a formula C_nH_{2n-2-x}Br_x, where "n" is 3 or greater and "x" is 2n-1 or less but not less than one. Thus, in general, they include all linear and branched hydrobromoalkenes having one or more carbon-carbon double bonds and having the general formula C_nH_{2n-2w+x}Br_x, where "w" is the number of carbon-carbon double bonds, "n" is w+1 or greater, and "x" is 2n-2w+1 or less but not less than one.

20 Hydrobromoarenes

Hydrobromoarenes include, by way of example only, the monobromo compounds bromobenzene (C₆H₅Br), bromonaphthalene (C₁₀H₇Br, 2 isomers), and bromobiphenyl (C₆H₅-C₆H₄Br, 3 isomers); and the dibromo compounds dibromobenzene (C₆H₄Br₂, 3 isomers),

20

dibromonaphthalene ($C_{10}H_6Br_2$, 6 isomers), and bromobiphenyl ($C_6H_5-C_6H_3Br_2$, 6 isomers, and $C_6H_4Br-C_6H_4Br$, 6 isomers); and brominated aromatics containing one or more hydrogen atoms and three or more bromine atoms.

5 Fluorine-Containing Component

10 A fluorine-containing component is added to the bromine-containing component to form the agent blends. The purpose of the fluorine-containing component is to produce an agent that resembles halons and HBFCs in fires. The fluorine-containing component may also aid to distribute the agent, modify the physical properties, reduce the toxicity, or to provide other benefits. The fluorine-containing component may be comprised of any organic compound containing fluorine or fluorine and chlorine but not containing any other halogen. Blends of the fluorine-containing component with hydrobromoalkanes, 15 hydrobromoalkenes, and/or hydrobromoarenes may be either azeotropes, which do not change in composition as they evaporate, or zeotropes, which do change in composition during evaporation (more volatile components tend to evaporate preferentially). Mixtures that change only slightly in composition during evaporation are sometimes termed "near azeotropes." In some cases, there are advantages to azeotropes and near azeotropes. Mixtures covered by this application include azeotropes, near azeotropes, and zeotropes.

20 The fluorine-containing component is comprised of non-brominated halocarbons. The halocarbons can be such materials as hydrochlorofluorocarbons, hydrofluorocarbons, perfluorocarbons, 25 perfluoroethers, hydrofluoroethers, hydrofluoropolyethers, and halogenated aromatics. Here, except for the aromatics, we use these terms to include both saturated and unsaturated hydrocarbons. Aromatics are always unsaturated. Hydrochlorofluorocarbons (HCFCs) are chemicals containing only hydrogen, chlorine, fluorine, 30 and carbon. Examples of HCFCs that could be incorporated into the fluorine-containing component are 2,2-dichloro-1,1,1-trifluoroethane

(CHCl₂CF₃), chlorodifluoromethane²¹ (CHClF₂), 2-chloro-1,1,1,2-tetrafluoroethane (CHClF₂CF₃), and 1-chloro-1,1-difluoroethane (CH₃CClF₂). Hydrofluorocarbons (HFCs) are chemicals containing only hydrogen, fluorine, and carbon. Examples of potential HFCs that could be incorporated into the fluorine-containing component are trifluoromethane (CHF₃), difluoromethane (CH₂F₂), 1,1-difluoroethane (CH₃CHF₂), pentafluoroethane (CHF₂CF₃), 1,1,1,2-tetrafluoroethane (CH₂FCF₃), 1,1,1,2,2-pentafluoropropane (CF₃CF₂CH₃), 1,1,1,2,3,3-hexafluoropropane (CF₃CHFCHF₂), 1,1,1,3,3,3-hexafluoropropane (CF₃CH₂CF₃), 1,1,1,2,2,3,3-heptafluoropropane (CF₃CF₂CF₂H), 1,1,1,2,3,3,3-heptafluoropropane (CF₃CHF₂CF₃), 1,1,1,4,4,4-hexafluorobutane (CF₃CH₂CH₂CF₃), and 1,1,1,2,2,3,4,5,5,5-decafluoropentane (CF₃CHFCH₂CF₂CF₃). Perfluorocarbons contain only fluorine and carbon. The saturated PFCs are characterized by very low toxicities. Examples of saturated perfluorocarbons that could be incorporated into the fluorine-containing component are tetrafluoromethane (CF₄), hexafluoroethane (CF₃CF₃), octafluoropropane (CF₃CF₂CF₃), decafluorobutane (CF₃CF₂CF₂CF₃), dodecafluoropentane (CF₃CF₂CF₂CF₂CF₃), tetradecafluorohexane (CF₃CF₂CF₂CF₂CF₂CF₃), perfluoromethylcyclohexane (C₆F₁₁CF₃), perfluorodimethylcyclohexane (C₆F₁₀(CF₃)₂), and perfluoromethyldecalin (C₁₀F₁₇CF₃). Examples of unsaturated perfluorocarbons that could be incorporated into the fluorine-containing component are perfluoro-1-butene (CF₂=CF₂CF₂CF₃) and perfluoro-2-butene (CF₃CF=CF₂CF₃). Perfluoroethers are compounds containing only carbon, oxygen, and fluorine and possessing an ether linkage (C-O-C). Examples are perfluorodimethyl ether (CF₃OCF₃), perfluoromethylethylether (CF₃CF₂OCF₃), perfluoromethylpropyl ether (CF₃OCF₂CF₂CF₃), and perfluorodiethyl ether (CF₃CF₂OCF₂CF₃). Hydrofluoroethers contain an ether linkage and the elements hydrogen, fluorine, carbon, and oxygen. Examples are methyl perfluorobutyl ether (CF₃CF₂CF₂CF₂OCH₃), ethyl perfluorobutyl ether

22

($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_2\text{OC}_2\text{H}_5$), bisdifluoromethyl ether ($\text{CHF}_2\text{OCHF}_2$), difluoromethyl 2,2,2-trifluoroethyl ether ($\text{CF}_3\text{CH}_2\text{OCHF}_2$), difluoromethyl 1,2,2,2-tetrafluoroethyl ether ($\text{CHF}_2\text{OCHF}_2\text{CF}_3$), methyl 1,1,2,2-tetrafluoroethyl ether ($\text{CH}_3\text{OCF}_2\text{CHF}_2$), methyl perfluoropropyl ether ($\text{CH}_3\text{OCF}_2\text{CF}_2\text{CF}_3$), methyl perfluoroisopropyl ether ($\text{CH}_3\text{OCF}(\text{CF}_3)_2$), 2,2,2-trifluoroethyl perfluoroethyl ether ($\text{CF}_3\text{CH}_2\text{OCF}_2\text{CF}_3$), and methyl perfluoroethyl ether ($\text{CH}_3\text{OCF}_2\text{CF}_3$). Hydrofluoropolyethers are polymeric liquids containing an ether linkage and the elements hydrogen, fluorine, carbon, and oxygen. Halogenated aromatics contain one or more 6-membered benzene rings. An example is chloropentafluorobenzene ($\text{C}_6\text{F}_5\text{Cl}$).

Description of Specific Embodiments

These and other aspects of the present invention will be more apparent upon consideration of the following examples.

As indicated previously, the present invention discloses the use of agents comprised of a bromine containing component and a fluorine containing component for the four applications of fire extinguishment or suppression using a total-flood application, fire extinguishment or suppression using a streaming application, explosion suppression, and inertion against fires and explosions. The bromine-containing component is comprised of one or more hydrobromocarbons selected from the group consisting of hydrobromoalkanes, hydrobromoalkenes, and hydrobromoarenes. The fluorine-containing component is comprised of one or more nonbrominated fluorine-containing halocarbons, which also contain no iodine. The following examples illustrate the fire and explosion protection in accordance with the invention.

Example 1. Into a flowing air stream in a cup burner apparatus in which a cup of burning *n*-heptane fuel was contained was introduced a mixture of 25 percent by weight 1-bromopropane ($\text{CH}_2\text{BrCH}_2\text{CH}_3$) and 75 percent by weight of a hydrofluoropolyether sufficient to raise the concentration of the blend in the air stream to 3.18 percent agent by

23

gas volume. A second test was run with a mixture of 10 percent by weight 1-bromopropane and 90 percent by weight of a hydrofluoropolyether sufficient to raise the concentration of the blend in the air stream to 3.11 percent agent by gas volume. Both mixtures extinguished the fire. These extinguishment concentrations exhibited by the blends in air were less than the extinguishment concentration of Halon 1211 in air (3.2 percent) and only slightly higher than the extinguishment concentration of Halon 1301 in air (2.9 percent) under the same conditions. A third test was run with 100 percent 1-bromopropane sufficient to raise the concentration of the agent in the air stream to 4.63 percent agent by gas volume. Thus, extinguishment by 1-bromopropane by itself required a concentration 46 percent greater than that of the blend in the first test and 49 percent greater than that of the blend in the second test showing the improvement achieved by the addition of the fluorine-containing component.

Example 2. Into a flowing air stream in a cup burner apparatus in which a cup of burning n-heptane fuel was contained was introduced a mixture of 11.5 percent by weight 2,3-dibromopentane ($\text{CH}_3\text{CHBrCHBrCH}_2\text{CH}_3$) and 88.5 percent by weight of 1,1,1,3,3,3-hexafluoropropane ($\text{CF}_3\text{CH}_2\text{CF}_3$) sufficient to raise the concentration of the blend in the air stream to 3.66 percent agent by gas volume. The mixture extinguished the flame. The extinguishment concentration of this blend was 46 percent less than the average extinguishment concentration (seven determinations) of 6.72 percent agent by gas volume found for 1,1,1,3,3,3-hexafluoropropane ($\text{CF}_3\text{CH}_2\text{CF}_3$) alone under the same conditions showing the improvement achieved by the addition of the bromine-containing component.

Example 3. Into a flowing air stream in a cup burner apparatus in which a cup of burning n-heptane fuel was contained was introduced a mixture of 11.4 percent by weight 2,3-dibromobutane ($\text{CH}_3\text{CHBrCHBrCH}_3$) and 88.6 percent by weight of 1,1,1,3,3,3-hexafluoropropane ($\text{CF}_3\text{CH}_2\text{CF}_3$) sufficient to raise the concentration of

24

the blend in the air stream to 4.64 percent agent by gas volume. The mixture extinguished the flame. The concentration in air required for extinguishment by this blend was 31 percent less than the average extinguishment concentration (seven determinations) of 6.72 percent agent by gas volume required to extinguish the fire with 1,1,1,3,3,3-hexafluoropropane ($\text{CF}_3\text{CH}_2\text{CF}_3$) alone.

Example 4. Onto a 2.25-square foot pan containing burning *n*-heptane fuel, a stream of a mixture of 25 percent by weight 1-bromopropane ($\text{CH}_2\text{BrCH}_2\text{CH}_3$) and 75 percent by weight of a hydrofluoropolyether was discharged using a flow rate of 0.29 pounds per second. The fire was extinguished in 2.6 seconds. In a second test using an identical apparatus, a mixture of 10 percent by weight 1-bromopropane and 90 percent by weight of a hydrofluoropolyether was discharged using a flow rate of 0.17 pounds per second. The fire was extinguished in 6 seconds.

Example 5. Onto a 2.25-square foot pan containing burning *n*-heptane fuel, a stream of a mixture of 25 percent by weight 1-bromopropane ($\text{CH}_2\text{BrCH}_2\text{CH}_3$) and 75 percent by weight of the hydrofluorocarbon 1,1,1,3,3,3-hexafluoropropane ($\text{CF}_3\text{CH}_2\text{CF}_3$) was discharged using a flow rate of 0.18 pounds per second. The fire was extinguished in 4.1 seconds.

Example 6. Into a well-ventilated 79.6-cubic foot test chamber containing an 8-inch diameter pan with a 1-inch deep pool of burning heptane was discharged 1.51 pounds of a blend of 15 percent 1-bromopropane ($\text{CH}_2\text{BrCH}_2\text{CH}_3$) and 85 percent by weight of a commercialized fire extinguishing agent NAF S-III, which is comprised of three HCFCs—chlorodifluoromethane (CHClF_2), 2-chloro-1,1,1,2-tetrafluoroethane ($\text{CHClF}_2\text{CF}_3$), and 2,2-dichloro-1,1,1-trifluoroethane (CHCl_2CF_3). The fire was extinguished in 5 seconds.

Example 7. An explosion occurs within a manufacturing facility used for filling aerosol cans with hydrocarbon propellants, and upon detection of the expanding fire ball, a blend of 20 percent by weight

25

3,3-dibromopropene ($\text{CH}_2=\text{CHCHBr}_2$) and 80 percent by weight decafluorobutane ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_3$) is automatically discharged and the explosion is suppressed.

5 Example 8. Upon detection of an unsafe concentration of methane in an enclosed room within a petroleum processing facility, a blend of 10 percent 1,1-dibromoethane (CHBr_2CH_3) and 90 percent perfluoro-1-butene ($\text{CF}_2=\text{CFCF}_2\text{CF}_3$) is discharged and the area is inerted to prevent an explosion or fire from occurring while the unsafe methane concentration condition is corrected.

10 The present invention has been described and illustrated with reference to certain preferred embodiments. Nevertheless, it will be understood that various modifications, alterations and substitutions may be apparent to one of ordinary skill in the art, and that such modifications, alterations and substitutions may be made without
15 departing from the essential invention. Thus, the present invention is, of course, in no way restricted to the specific disclosure of the specification and examples, but also encompasses any modifications within the scope of the appended claims.

We claim: 26

1. A method of extinguishing or suppressing a fire in a total-flood application, said method comprising the steps of:

- a) providing at least one hydrobromocarbon selected from the group consisting of hydrobromoalkanes,
hydrobromoalkenes, and hydrobromoarenes,
- b) mixing said at least one hydrobromocarbon with at least one halocarbon selected from the group consisting of all fluorine-containing nonbrominated halocarbons to give a blend, said halocarbons containing no bromine and no iodine.
- c) disposing said blend in a pressurized discharge system, and
- d) discharging said blend into an area to extinguish or suppress fires in that area.

2. The method of claim 1 wherein said at least one halocarbon is selected from the group consisting of saturated and unsaturated hydrochlorofluorocarbons, hydrofluorocarbons, perfluorocarbons, perfluoroethers, hydrofluoroethers, hydrofluoropolyethers and halogenated aromatics, said hydrobromoalkanes either have the formula $C_nH_{2n+2-x}Br_x$, where n is 2 or greater and x is at least 1 but not greater than 2n+1, or have the formula $C_nH_{2n+2-2y-x}Br_x$, where n is 3 or greater, x is at least 1 but not greater than 2n+1-2y, and y is the number of rings, and said hydrobromoalkenes have the formula $C_nH_{2n-2w+2-x}Br_x$, where w is the number of carbon-carbon double bonds, n is w+1 or greater, and x is 2n-2w+1 or less but at least 1.

3. The method of claim 2 wherein said at least one hydrobromocarbon is selected from the group consisting of bromoethane (CH_3CH_2Br), 1-bromopropane ($CH_3CH_2CH_2Br$), 2-bromopropane ($CH_3CHBrCH_3$), 1-bromobutane ($CH_3CH_2CH_2CH_2Br$), 2-bromobutane ($CH_3CH_2CHBrCH_3$), 1-bromopentane ($CH_3CH_2CH_2CH_2CH_2Br$), 2-bromopentane ($CH_3CH_2CH_2CHBrCH_3$), 3-bromopentane ($CH_3CH_2CHBrCH_2CH_3$), 1-bromohexane ($CH_3CH_2CH_2CH_2CH_2CH_2Br$), 2-bromohexane

- (CH₃CH₂CH₂CH₂CHBrCH₃), 3-bromohexane
- 10 (CH₃CH₂CHBrCH₂CH₂CH₃), 1-bromo-2-methylpropane
- (CH₃CH(CH₃)CH₂Br), 2-bromo-2-methylpropane (CH₃C(CH₃)BrCH₃),
- 1-bromo-2-methylbutane (CH₃CH₂CH(CH₃)CH₂Br), 1-bromo-3-
- methylbutane (CH₃CH(CH₃)CH₂CH₂Br), 1-bromo-2-methylpentane
- (CH₃CH₂CH₂CH(CH₃)CH₂Br), 1-bromo-3-methylpentane
- 15 (CH₃CH₂CH(CH₃)CH₂CH₂Br), 1-bromo-4-methylpentane
- (CH₃CH(CH₃)CH₂CH₂CH₂Br), 1,1-dibromoethane (CH₃CHBr₂), 1,2-
- dibromoethane (CH₂BrCH₂Br), 1,1-dibromopropane (CH₃CH₂CHBr₂),
- 1,2-dibromopropane (CH₃CHBrCH₂Br), 1,3-dibromopropane
- (CH₂BrCH₂CH₂Br), bromocyclopropane (C₃H₅Br), bromocyclobutane
- 20 (C₄H₇Br), bromocyclopentane (C₅H₉Br), bromobenzene (C₆H₅Br),
- dibromobenzene (C₆H₄Br₂), bromotoluene (C₇H₇Br), 3,3-
- dibromopropene (CH₂=CHCHBr₂), bromoethene (CH₂=CHBr), 2-
- methyl-1,1,3-tribromopropene (CBr₂=C(CH₃)CH₂Br), 3-bromopropene
- (CH₂=CHCH₂Br), 2-bromopropene (CH₂=CBrCH₃), 1-bromopropene
- 25 (CHBr=CHCH₃), 2,3-dibromopropene (CH₂=CBrCH₂Br), 2-methyl-1-
- bromopropene (CHBr=C(CH₃)CH₃), 4-bromo-1,3-butanediene
- (CH₂=CHCH=CHBr), 3-bromo-1,3-butanediene (CH₂=CHCBr=CH₂),
- 3-bromo-2-methyl-1,3-butanediene (CH₂=C(CH₃)CBr=CH₂), and 3,4-
- dibromo-2-methyl-1,3-butanediene (CH₂=C(CH₃)CBr=CHBr) and
- 30 isomers thereof and said at least one halocarbon is selected from the
- group consisting of 2,2-dichloro-1,1,1-trifluoroethane (CHCl₂CF₃),
- chlorodifluoromethane (CHClF₂), 2-chloro-1,1,1,2-tetrafluoroethane
- (CHClF₂CF₃), 1-chloro-1,1-difluoroethane (CH₃CClF₂),
- trifluoromethane (CHF₃), difluoromethane (CH₂F₂), 1,1-difluoroethane
- 35 (CH₃CHF₂), pentafluoroethane (CHF₂CF₃), 1,1,1,2-tetrafluoroethane
- (CH₂FCF₃), 1,1,1,2,2-pentafluoropropane (CF₃CF₂CH₃), 1,1,1,2,3,3-
- hexafluoropropane (CF₃CHFCHF₂), 1,1,1,3,3,3-hexafluoropropane
- (CF₃CH₂CF₃), 1,1,1,2,2,3,3-heptafluoropropane (CF₃CF₂CF₂H),
- 1,1,1,2,3,3,3-heptafluoropropane (CF₃CHF₂CF₃), 1,1,1,4,4,4-
- 40 hexafluorobutane (CF₃CH₂CH₂CF₃), 1,1,1,2,2,3,4,5,5,5-

decafluoropentane ($\text{CF}_3\text{CHFCHF}\text{CF}_2\text{CF}_3$), tetrafluoromethane (CF_4),
 hexafluoroethane (CF_3CF_3), octafluoropropane ($\text{CF}_3\text{CF}_2\text{CF}_3$),
 decafluorobutane ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_3$), dodecafluoropentane
 ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_2\text{CF}_3$), tetradecafluorohexane ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_2\text{CF}_2\text{CF}_3$),
 45 perfluoromethylcyclohexane ($\text{C}_6\text{F}_{11}\text{CF}_3$), perfluorodimethylcyclohexane
 ($\text{C}_6\text{F}_{10}(\text{CF}_3)_2$), perfluoromethyldecalin ($\text{C}_{10}\text{F}_{17}\text{CF}_3$), perfluorodimethyl
 ether (CF_3OCF_3), perfluoromethylethylether ($\text{CF}_3\text{CF}_2\text{OCF}_3$),
 perfluoromethylpropyl ether ($\text{CF}_3\text{OCF}_2\text{CF}_2\text{CF}_3$), perfluorodiethyl ether
 ($\text{CF}_3\text{CF}_2\text{OCF}_2\text{CF}_3$), methyl perfluorobutyl ether ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_2\text{OCH}_3$),
 50 ethyl perfluorobutyl ether ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_2\text{OC}_2\text{H}_5$), bisdifluoromethyl
 ether ($\text{CHF}_2\text{OCHF}_2$), difluoromethyl 2,2,2-trifluoroethyl ether
 ($\text{CF}_3\text{CH}_2\text{OCHF}_2$), difluoromethyl 1,2,2,2-tetrafluoroethyl ether
 ($\text{CHF}_2\text{OCHF}\text{CF}_3$), methyl 1,1,2,2-tetrafluoroethyl ether
 ($\text{CH}_3\text{OCF}_2\text{CHF}_2$), methyl perfluoropropyl ether ($\text{CH}_3\text{OCF}_2\text{CF}_2\text{CF}_3$),
 55 methyl perfluoroisopropyl ether ($\text{CH}_3\text{OCF}(\text{CF}_3)_2$), 2,2,2-trifluoroethyl
 perfluoroethyl ether ($\text{CF}_3\text{CH}_2\text{OCF}_2\text{CF}_3$), methyl perfluoroethyl ether
 ($\text{CH}_3\text{OCF}_2\text{CF}_3$), perfluoro-1-butene ($\text{CF}_2=\text{CF}\text{CF}_2\text{CF}_3$), perfluoro-2-
 butene ($\text{CF}_3\text{CF}=\text{CF}\text{CF}_3$), and chloropentafluorobenzene ($\text{C}_6\text{F}_5\text{Cl}$).

4. A method of extinguishing or suppressing a fire in a
 streaming application, said method comprising the steps of

- a) providing at least one hydrobromocarbon selected from the
 group consisting of hydrobromoalkanes,
 5 hydrobromoalkenes, and hydrobromoarenes,
- b) mixing said at least one hydrobromocarbon with at least one
 halocarbon selected from the group consisting of all
 fluorine-containing halocarbons to give a blend, said
 halocarbons containing no bromine and no iodine.
- 10 c) disposing said blend in a pressurized discharge system, and
- d) discharging said blend from said system toward an existing
 fire to suppress or extinguish said fire.

5. The method of claim 4 wherein said at least one halocarbon
 is selected from the group consisting of saturated and unsaturated
 hydrochlorofluorocarbons, hydrofluorocarbons, perfluorocarbons,

perfluoroethers, hydrofluoroethers, hydrofluoropolyethers and
 5 halogenated aromatics, said hydrobromoalkanes either have the formula
 $C_nH_{2n+2-x}Br_x$, where n is 2 or greater and x is at least 1 but not greater
 than $2n+1$, or have the formula $C_nH_{2n+2-2y-x}Br_x$, where n is 3 or greater,
 x is at least 1 but not greater than $2n+1-2y$, and y is the number of
 10 rings, and said hydrobromoalkenes have the formula $C_nH_{2n-2w+2-x}Br_x$,
 where w is the number of carbon-carbon double bonds, n is w+1 or
 greater, and x is $2n-2w+1$ or less but at least 1.

6. The method of claim 5 wherein said at least one
 hydrobromocarbon is selected from the group consisting of
 bromoethane (CH_3CH_2Br), 1-bromopropane ($CH_3CH_2CH_2Br$), 2-
 bromopropane ($CH_3CHBrCH_3$), 1-bromobutane ($CH_3CH_2CH_2CH_2Br$),
 5 2-bromobutane ($CH_3CH_2CHBrCH_3$), 1-bromopentane
 ($CH_3CH_2CH_2CH_2CH_2Br$), 2-bromopentane ($CH_3CH_2CH_2CHBrCH_3$),
 3-bromopentane ($CH_3CH_2CHBrCH_2CH_3$), 1-bromohexane
 ($CH_3CH_2CH_2CH_2CH_2CH_2Br$), 2-bromohexane
 ($CH_3CH_2CH_2CH_2CHBrCH_3$), 3-bromohexane
 10 ($CH_3CH_2CHBrCH_2CH_2CH_3$), 1-bromo-2-methylpropane
 ($CH_3CH(CH_3)CH_2Br$), 2-bromo-2-methylpropane ($CH_3C(CH_3)BrCH_3$),
 1-bromo-2-methylbutane ($CH_3CH_2CH(CH_3)CH_2Br$), 1-bromo-3-
 methylbutane ($CH_3CH(CH_3)CH_2CH_2Br$), 1-bromo-2-methylpentane
 ($CH_3CH_2CH_2CH(CH_3)CH_2Br$), 1-bromo-3-methylpentane
 15 ($CH_3CH_2CH(CH_3)CH_2CH_2Br$), 1-bromo-4-methylpentane
 ($CH_3CH(CH_3)CH_2CH_2CH_2Br$), 1,1-dibromoethane (CH_3CHBr_2), 1,2-
 dibromoethane (CH_2BrCH_2Br), 1,1-dibromopropane ($CH_3CH_2CHBr_2$),
 1,2-dibromopropane ($CH_3CHBrCH_2Br$), 1,3-dibromopropane
 ($CH_2BrCH_2CH_2Br$), bromocyclopropane (C_3H_5Br), bromocyclobutane
 20 (C_4H_7Br), bromocyclopentane (C_5H_9Br), bromobenzene (C_6H_5Br),
 dibromobenzene ($C_6H_4Br_2$), bromotoluene (C_7H_7Br), 3,3-
 dibromopropene ($CH_2=CHCHBr_2$), bromoethene ($CH_2=CHBr$), 2-
 methyl-1,1,3-tribromopropene ($CBr_2=C(CH_3)CH_2Br$), 3-bromopropene
 ($CH_2=CHCH_2Br$), 2-bromopropene ($CH_2=CBrCH_3$), 1-bromopropene
 25 ($CHBr=CHCH_3$), 2,3-dibromopropene ($CH_2=CBrCH_2Br$), 2-methyl-1-

bromopropene ($\text{CHBr}=\text{C}(\text{CH}_3)\text{CH}_3$), 4-bromo-1,3-butanediene
 ($\text{CH}_2=\text{CHCH}=\text{CHBr}$), 3-bromo-1,3-butanediene ($\text{CH}_2=\text{CHCBr}=\text{CH}_2$),
 3-bromo-2-methyl-1,3-butanediene ($\text{CH}_2=\text{C}(\text{CH}_3)\text{CBr}=\text{CH}_2$), and 3,4-
 dibromo-2-methyl-1,3-butanediene ($\text{CH}_2=\text{C}(\text{CH}_3)\text{CBr}=\text{CHBr}$) and
 isomers thereof. and said at least one halocarbon is selected from the
 group consisting of 2,2-dichloro-1,1,1-trifluoroethane (CHCl_2CF_3),
 chlorodifluoromethane (CHClF_2), 2-chloro-1,1,1,2-tetrafluoroethane
 ($\text{CHClF}_2\text{CF}_3$), 1-chloro-1,1-difluoroethane (CH_3CClF_2),
 trifluoromethane (CHF_3), difluoromethane (CH_2F_2), 1,1-difluoroethane
 (CH_3CHF_2), pentafluoroethane (CHF_2CF_3), 1,1,1,2-tetrafluoroethane
 ($\text{CH}_2\text{F}_2\text{CF}_3$), 1,1,1,2,2-pentafluoropropane ($\text{CF}_3\text{CF}_2\text{CH}_3$), 1,1,1,2,3,3-
 hexafluoropropane ($\text{CF}_3\text{CHFCHF}_2$), 1,1,1,3,3,3-hexafluoropropane
 ($\text{CF}_3\text{CH}_2\text{CF}_3$), 1,1,1,2,2,3,3-heptafluoropropane ($\text{CF}_3\text{CF}_2\text{CF}_2\text{H}$),
 1,1,1,2,3,3,3-heptafluoropropane ($\text{CF}_3\text{CHF}_2\text{CF}_3$), 1,1,1,4,4,4-
 hexafluorobutane ($\text{CF}_3\text{CH}_2\text{CH}_2\text{CF}_3$), 1,1,1,2,2,3,4,5,5,5-
 decafluoropentane ($\text{CF}_3\text{CHFCH}_2\text{CF}_2\text{CF}_3$), tetrafluoromethane (CF_4),
 hexafluoroethane (CF_3CF_3), octafluoropropane ($\text{CF}_3\text{CF}_2\text{CF}_3$),
 decafluorobutane ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_3$), dodecafluoropentane
 ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_2\text{CF}_3$), tetradecafluorohexane ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_2\text{CF}_2\text{CF}_3$),
 perfluoromethylcyclohexane ($\text{C}_6\text{F}_{11}\text{CF}_3$), perfluorodimethylcyclohexane
 ($\text{C}_6\text{F}_{10}(\text{CF}_3)_2$), perfluoromethyldecalin ($\text{C}_{10}\text{F}_{17}\text{CF}_3$), perfluorodimethyl
 ether (CF_3OCF_3), perfluoromethylethylether ($\text{CF}_3\text{CF}_2\text{OCF}_3$),
 perfluoromethylpropyl ether ($\text{CF}_3\text{OCF}_2\text{CF}_2\text{CF}_3$), perfluorodiethyl ether
 ($\text{CF}_3\text{CF}_2\text{OCF}_2\text{CF}_3$), methyl perfluorobutyl ether ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_2\text{OCH}_3$),
 ethyl perfluorobutyl ether ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_2\text{OC}_2\text{H}_5$), bisdifluoromethyl
 ether ($\text{CHF}_2\text{OCHF}_2$), difluoromethyl 2,2,2-trifluoroethyl ether
 ($\text{CF}_3\text{CH}_2\text{OCHF}_2$), difluoromethyl 1,2,2,2-tetrafluoroethyl ether
 ($\text{CHF}_2\text{OCHF}_2\text{CF}_3$), methyl 1,1,2,2-tetrafluoroethyl ether
 ($\text{CH}_3\text{OCF}_2\text{CHF}_2$), methyl perfluoropropyl ether ($\text{CH}_3\text{OCF}_2\text{CF}_2\text{CF}_3$),
 methyl perfluoroisopropyl ether ($\text{CH}_3\text{OCF}(\text{CF}_3)_2$), 2,2,2-trifluoroethyl
 perfluoroethyl ether ($\text{CF}_3\text{CH}_2\text{OCF}_2\text{CF}_3$), methyl perfluoroethyl ether

31

(CH₃OCF₂CF₃), perfluoro-1-butene (CF₂=CF CF₂CF₃), perfluoro-2-butene (CF₃CF=CF CF₃), and chloropentafluorobenzene (C₆F₅Cl).

7. A method of suppressing an explosion, said method comprising the steps of

- a) providing at least one hydrobromocarbon selected from the group consisting of hydrobromoalkanes,
hydrobromoalkenes, and hydrobromoarenes,
- b) mixing said at least one hydrobromocarbon with at least one halocarbon selected from the group consisting of all fluorine-containing halocarbons to give a blend, said halocarbons containing no bromine and no iodine,
- c) detecting an explosion and discharging said blend into the area of the explosion to suppress the explosion.

8. The method of claim 7 wherein said at least one halocarbon is selected from the group consisting of saturated and unsaturated hydrochlorofluorocarbons, hydrofluorocarbons, perfluorocarbons, perfluoroethers, hydrofluoroethers, hydrofluoropolyethers and halogenated aromatics, said hydrobromoalkanes either have the formula C_nH_{2n+2-x}Br_x, where n is 2 or greater and x is at least 1 but not greater than 2n+1, or have the formula C_nH_{2n+2-2y-x}Br_x, where n is 3 or greater, x is at least 1 but not greater than 2n+1-2y, and y is the number of rings, and said hydrobromoalkenes have the formula C_nH_{2n-2w+2-x}Br_x, where w is the number of carbon-carbon double bonds, n is w+1 or greater, and x is 2n-2w+1 or less but at least 1.

9. The method of claim 8 wherein said at least one hydrobromocarbon is selected from the group consisting of bromoethane (CH₃CH₂Br), 1-bromopropane (CH₃CH₂CH₂Br), 2-bromopropane (CH₃CHBrCH₃), 1-bromobutane (CH₃CH₂CH₂CH₂Br), 2-bromobutane (CH₃CH₂CHBrCH₃), 1-bromopentane (CH₃CH₂CH₂CH₂CH₂Br), 2-bromopentane (CH₃CH₂CH₂CHBrCH₃), 3-bromopentane (CH₃CH₂CHBrCH₂CH₃), 1-bromohexane (CH₃CH₂CH₂CH₂CH₂CH₂Br), 2-bromohexane (CH₃CH₂CH₂CH₂CHBrCH₃), 3-bromohexane (CH₃CH₂CH₂CHBrCH₂CH₃).

- 10 $(\text{CH}_3\text{CH}_2\text{CHBrCH}_2\text{CH}_2\text{CH}_3)$, 1-bromo-2-methylpropane
 $(\text{CH}_3\text{CH}(\text{CH}_3)\text{CH}_2\text{Br})$, 2-bromo-2-methylpropane $(\text{CH}_3\text{C}(\text{CH}_3)\text{BrCH}_3)$,
 1-bromo-2-methylbutane $(\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{Br})$, 1-bromo-3-
 methylbutane $(\text{CH}_3\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{Br})$, 1-bromo-2-methylpentane
 $(\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{Br})$, 1-bromo-3-methylpentane
 15 $(\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{Br})$, 1-bromo-4-methylpentane
 $(\text{CH}_3\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_2\text{Br})$, 1,1-dibromoethane $(\text{CH}_3\text{CHBr}_2)$, 1,2-
 dibromoethane $(\text{CH}_2\text{BrCH}_2\text{Br})$, 1,1-dibromopropane $(\text{CH}_3\text{CH}_2\text{CHBr}_2)$,
 1,2-dibromopropane $(\text{CH}_3\text{CHBrCH}_2\text{Br})$, 1,3-dibromopropane
 $(\text{CH}_2\text{BrCH}_2\text{CH}_2\text{Br})$, bromocyclopropane $(\text{C}_3\text{H}_5\text{Br})$, bromocyclobutane
 20 $(\text{C}_4\text{H}_7\text{Br})$, bromocyclopentane $(\text{C}_5\text{H}_9\text{Br})$, bromobenzene $(\text{C}_6\text{H}_5\text{Br})$,
 dibromobenzene $(\text{C}_6\text{H}_4\text{Br}_2)$, bromotoluene $(\text{C}_7\text{H}_7\text{Br})$, 3,3-
 dibromopropene $(\text{CH}_2=\text{CHCHBr}_2)$, bromoethene $(\text{CH}_2=\text{CHBr})$, 2-
 methyl-1,1,3-tribromopropene $(\text{CBr}_2=\text{C}(\text{CH}_3)\text{CH}_2\text{Br})$, 3-bromopropene
 $(\text{CH}_2=\text{CHCH}_2\text{Br})$, 2-bromopropene $(\text{CH}_2=\text{CBrCH}_3)$, 1-bromopropene
 25 $(\text{CHBr}=\text{CHCH}_3)$, 2,3-dibromopropene $(\text{CH}_2=\text{CBrCH}_2\text{Br})$, 2-methyl-1-
 bromopropene $(\text{CHBr}=\text{C}(\text{CH}_3)\text{CH}_3)$, 4-bromo-1,3-butanediene
 $(\text{CH}_2=\text{CHCH}=\text{CHBr})$, 3-bromo-1,3-butanediene $(\text{CH}_2=\text{CHCBr}=\text{CH}_2)$,
 3-bromo-2-methyl-1,3-butanediene $(\text{CH}_2=\text{C}(\text{CH}_3)\text{CBr}=\text{CH}_2)$, and 3,4-
 dibromo-2-methyl-1,3-butanediene $(\text{CH}_2=\text{C}(\text{CH}_3)\text{CBr}=\text{CHBr})$ and
 30 isomers thereof. and said at least one halocarbon is selected from the
 group consisting of 2,2-dichloro-1,1,1-trifluoroethane $(\text{CHCl}_2\text{CF}_3)$,
 chlorodifluoromethane (CHClF_2) , 2-chloro-1,1,1,2-tetrafluoroethane
 $(\text{CHClF}_2\text{CF}_3)$, 1-chloro-1,1-difluoroethane $(\text{CH}_3\text{CClF}_2)$,
 trifluoromethane (CHF_3) , difluoromethane (CH_2F_2) , 1,1-difluoroethane
 35 $(\text{CH}_3\text{CHF}_2)$, pentafluoroethane $(\text{CHF}_2\text{CF}_3)$, 1,1,1,2-tetrafluoroethane
 $(\text{CH}_2\text{FCF}_3)$, 1,1,1,2,2-pentafluoropropane $(\text{CF}_3\text{CF}_2\text{CH}_3)$, 1,1,1,2,3,3-
 hexafluoropropane $(\text{CF}_3\text{CHFCHF}_2)$, 1,1,1,3,3,3-hexafluoropropane
 $(\text{CF}_3\text{CH}_2\text{CF}_3)$, 1,1,1,2,2,3,3-heptafluoropropane $(\text{CF}_3\text{CF}_2\text{CF}_2\text{H})$,
 1,1,1,2,3,3,3-heptafluoropropane $(\text{CF}_3\text{CHF}_2\text{CF}_3)$, 1,1,1,4,4,4-
 40 hexafluorobutane $(\text{CF}_3\text{CH}_2\text{CH}_2\text{CF}_3)$, 1,1,1,2,2,3,4,5,5,5-
 decafluoropentane $(\text{CF}_3\text{CHFCH}_2\text{CF}_2\text{CF}_3)$, tetrafluoromethane (CF_4) ,

33

hexafluoroethane (CF_3CF_3), octafluoropropane ($\text{CF}_3\text{CF}_2\text{CF}_3$),
 decafluorobutane ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_3$), dodecafluoropentane
 ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_2\text{CF}_3$), tetradecafluorohexane ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_2\text{CF}_2\text{CF}_3$),
 45 perfluoromethylcyclohexane ($\text{C}_6\text{F}_{11}\text{CF}_3$), perfluorodimethylcyclohexane
 ($\text{C}_6\text{F}_{10}(\text{CF}_3)_2$), perfluoromethyldecalin ($\text{C}_{10}\text{F}_{17}\text{CF}_3$), perfluorodimethyl
 ether (CF_3OCF_3), perfluoromethylethylether ($\text{CF}_3\text{CF}_2\text{OCF}_3$),
 perfluoromethylpropyl ether ($\text{CF}_3\text{OCF}_2\text{CF}_2\text{CF}_3$), perfluorodiethyl ether
 ($\text{CF}_3\text{CF}_2\text{OCF}_2\text{CF}_3$), methyl perfluorobutyl ether ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_2\text{OCH}_3$),
 50 ethyl perfluorobutyl ether ($\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_2\text{OC}_2\text{H}_5$), bisdifluoromethyl
 ether ($\text{CHF}_2\text{OCHF}_2$), difluoromethyl 2,2,2-trifluoroethyl ether
 ($\text{CF}_3\text{CH}_2\text{OCHF}_2$), difluoromethyl 1,2,2,2-tetrafluoroethyl ether
 ($\text{CHF}_2\text{OCHF}_2\text{CF}_3$), methyl 1,1,2,2-tetrafluoroethyl ether
 ($\text{CH}_3\text{OCF}_2\text{CHF}_2$), methyl perfluoropropyl ether ($\text{CH}_3\text{OCF}_2\text{CF}_2\text{CF}_3$),
 55 methyl perfluoroisopropyl ether ($\text{CH}_3\text{OCF}(\text{CF}_3)_2$), 2,2,2-trifluoroethyl
 perfluoroethyl ether ($\text{CF}_3\text{CH}_2\text{OCF}_2\text{CF}_3$), methyl perfluoroethyl ether
 ($\text{CH}_3\text{OCF}_2\text{CF}_3$), perfluoro-1-butene ($\text{CF}_2=\text{CFCF}_2\text{CF}_3$), perfluoro-2-
 butene ($\text{CF}_3\text{CF}=\text{CFCF}_3$), and chloropentafluorobenzene ($\text{C}_6\text{F}_5\text{Cl}$).

10. A method of inerting an area to prevent a fire or explosion,
 said method comprising the steps of:

- a) providing at least one hydrobromocarbon selected from the
 group consisting of hydrobromoalkanes,
 5 hydrobromoalkenes, and hydrobromoarenes,
- b) mixing said at least one hydrobromocarbon with at least one
 halocarbon selected from the group consisting of all
 fluorine-containing halocarbons to give a blend, said
 halocarbon containing no bromine and no iodine,
- 10 c) disposing said blend in a pressurized discharge system, and
- d) discharging said agent into said area to prevent a fire or an
 explosion from occurring.

11. The method of claim 10 wherein said at least one halocarbon
 is selected from the group consisting of saturated and unsaturated
 hydrochlorofluorocarbons, hydrofluorocarbons, perfluorocarbons,
 perfluoroethers, hydrofluoroethers, hydrofluoropolyethers and

- 5 halogenated aromatics, said hydrobromoalkanes either have the formula $C_nH_{2n+2-x}Br_x$, where n is 2 or greater and x is at least 1 but not greater than 2n+1, or have the formula $C_nH_{2n+2-2y-x}Br_x$, where n is 3 or greater, x is at least 1 but not greater than 2n+1-2y, and y is the number of rings, and said hydrobromoalkenes have the formula $C_nH_{2n-2w+2-x}Br_x$,
 10 where w is the number of carbon-carbon double bonds, n is w+1 or greater, and x is 2n-2w+1 or less but at least 1.

12. The method of claim 11 wherein said at least one hydrobromocarbon is selected from the group consisting of bromoethane (CH_3CH_2Br), 1-bromopropane ($CH_3CH_2CH_2Br$), 2-bromopropane ($CH_3CHBrCH_3$), 1-bromobutane ($CH_3CH_2CH_2CH_2Br$),
 5 2-bromobutane ($CH_3CH_2CHBrCH_3$), 1-bromopentane ($CH_3CH_2CH_2CH_2CH_2Br$), 2-bromopentane ($CH_3CH_2CH_2CHBrCH_3$), 3-bromopentane ($CH_3CH_2CHBrCH_2CH_3$), 1-bromohexane ($CH_3CH_2CH_2CH_2CH_2CH_2Br$), 2-bromohexane ($CH_3CH_2CH_2CH_2CHBrCH_3$), 3-bromohexane ($CH_3CH_2CHBrCH_2CH_2CH_3$), 1-bromo-2-methylpropane ($CH_3CH(CH_3)CH_2Br$), 2-bromo-2-methylpropane ($CH_3C(CH_3)BrCH_3$), 1-bromo-2-methylbutane ($CH_3CH_2CH(CH_3)CH_2Br$), 1-bromo-3-methylbutane ($CH_3CH(CH_3)CH_2CH_2Br$), 1-bromo-2-methylpentane ($CH_3CH_2CH_2CH(CH_3)CH_2Br$), 1-bromo-3-methylpentane ($CH_3CH_2CH(CH_3)CH_2CH_2Br$), 1-bromo-4-methylpentane ($CH_3CH(CH_3)CH_2CH_2CH_2Br$), 1,1-dibromoethane (CH_3CHBr_2), 1,2-dibromoethane (CH_2BrCH_2Br), 1,1-dibromopropane ($CH_3CH_2CHBr_2$), 1,2-dibromopropane ($CH_3CHBrCH_2Br$), 1,3-dibromopropane ($CH_2BrCH_2CH_2Br$), bromocyclopropane (C_3H_5Br), bromocyclobutane (C_4H_7Br), bromocyclopentane (C_5H_9Br), bromobenzene (C_6H_5Br), dibromobenzene ($C_6H_4Br_2$), bromotoluene (C_7H_7Br), 3,3-dibromopropene ($CH_2=CHCHBr_2$), bromoethene ($CH_2=CHBr$), 2-methyl-1,1,3-tribromopropene ($CBr_2=C(CH_3)CH_2Br$), 3-bromopropene ($CH_2=CHCH_2Br$), 2-bromopropene ($CH_2=CBrCH_3$), 1-bromopropene ($CHBr=CHCH_3$), 2,3-dibromopropene ($CH_2=CBrCH_2Br$), 2-methyl-1-bromopropene ($CHBr=C(CH_3)CH_3$), 4-bromo-1,3-butanediene

35

(CH₂=CHCH=CHBr), 3-bromo-1,3-butanediene (CH₂=CHCBr=CH₂),
 3-bromo-2-methyl-1,3-butanediene (CH₂=C(CH₃)CBr=CH₂), and 3,4-
 dibromo-2-methyl-1,3-butanediene (CH₂=C(CH₃)CBr=CHBr) and
 30 isomers thereof. and said at least one halocarbon is selected from the
 group consisting of 2,2-dichloro-1,1,1-trifluoroethane (CHCl₂CF₃),
 chlorodifluoromethane (CHClF₂), 2-chloro-1,1,1,2-tetrafluoroethane
 (CHClF₂CF₃), 1-chloro-1,1-difluoroethane (CH₃CClF₂),
 trifluoromethane (CHF₃), difluoromethane (CH₂F₂), 1,1-difluoroethane
 35 (CH₃CHF₂), pentafluoroethane (CHF₂CF₃), 1,1,1,2-tetrafluoroethane
 (CH₂FCF₃), 1,1,1,2,2-pentafluoropropane (CF₃CF₂CH₃), 1,1,1,2,3,3-
 hexafluoropropane (CF₃CHFCHF₂), 1,1,1,3,3,3-hexafluoropropane
 (CF₃CH₂CF₃), 1,1,1,2,2,3,3-heptafluoropropane (CF₃CF₂CF₂H),
 1,1,1,2,3,3,3-heptafluoropropane (CF₃CHF₂CF₃), 1,1,1,4,4,4-
 40 hexafluorobutane (CF₃CH₂CH₂CF₃), 1,1,1,2,2,3,4,5,5,5-
 decafluoropentane (CF₃CHFCHF₂CF₂CF₃), tetrafluoromethane (CF₄),
 hexafluoroethane (CF₃CF₃), octafluoropropane (CF₃CF₂CF₃),
 decafluorobutane (CF₃CF₂CF₂CF₃), dodecafluoropentane
 (CF₃CF₂CF₂CF₂CF₃), tetradecafluorohexane (CF₃CF₂CF₂CF₂CF₂CF₃),
 45 perfluoromethylcyclohexane (C₆F₁₁CF₃), perfluorodimethylcyclohexane
 (C₆F₁₀(CF₃)₂), perfluoromethyldecalin (C₁₀F₁₇CF₃), perfluorodimethyl
 ether (CF₃OCF₃), perfluoromethylethylether (CF₃CF₂OCF₃),
 perfluoromethylpropyl ether (CF₃OCF₂CF₂CF₃), perfluorodiethyl ether
 (CF₃CF₂OCF₂CF₃), methyl perfluorobutyl ether (CF₃CF₂CF₂CF₂OCH₃),
 50 ethyl perfluorobutyl ether (CF₃CF₂CF₂CF₂OC₂H₅), bisdifluoromethyl
 ether (CHF₂OCHF₂), difluoromethyl 2,2,2-trifluoroethyl ether
 (CF₃CH₂OCHF₂), difluoromethyl 1,2,2,2-tetrafluoroethyl ether
 (CHF₂OCHF₂CF₃), methyl 1,1,2,2-tetrafluoroethyl ether
 (CH₃OCF₂CHF₂), methyl perfluoropropyl ether (CH₃OCF₂CF₂CF₃),
 55 methyl perfluoroisopropyl ether (CH₃OCF(CF₃)₂), 2,2,2-trifluoroethyl
 perfluoroethyl ether (CF₃CH₂OCF₂CF₃), methyl perfluoroethyl ether
 (CH₃OCF₂CF₃), perfluoro-1-butene (CF₂=CF₂CF₂CF₃), perfluoro-2-
 butene (CF₃CF=CF₂CF₃), and chloropentafluorobenzene (C₆F₅Cl).

1/1




Figure 1. Atmospheric Lifetime as a Function of Number of Hydrogen Atoms.